



ClariFly® Fly Control Block

For Cattle and Horses

Prevents the emergence of house flies, horn flies, face flies and stable flies from manure of treated cattle. Prevents the emergence of house flies and stable flies from manure of treated horses.

ACTIVE INGREDIENT:

Diflubenzuron (CAS #35367-38-5) 0.02%

OTHER INGREDIENTS 99.98%

TOTAL 100.00%

Contains 90.8 mg of Diflubenzuron per pound.

GUARANTEED ANALYSIS

Crude Protein (Min) 8.00%

Crude Fat (Min) 1.00%

Crude Fiber (Max) 4.00%

Calcium (Min) 11.00%

Calcium (Max) 13.20%

Phosphorus (Min) 1.00%

Salt (Min) 18.50%

Salt (Max) 22.20%

Cobalt (Min) 5 ppm

Copper (Min) 500 ppm

Iodine (Min) 25 ppm

Manganese (Min) 2,000 ppm

Selenium (Min) 13 ppm

Zinc (Min) 1,500 ppm

Vitamin A (Min) 50,000 IU/lb

Vitamin D-3 (Min) 5,000 IU/lb

Vitamin E (Min) 5 IU/lb

FEED INGREDIENTS

Processed Grain By-Products, Calcium Carbonate, Salt, Molasses Products, Calcium Hydroxide, Monocalcium Phosphate, Dicalcium Phosphate, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Selenium Yeast, Brewer's Dried Yeast, Ethylenediamine Dihydrochloride, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

KEEP OUT OF REACH OF CHILDREN

CAUTION

ClariFly® is a registered trademark of Wellmark International

Manufactured By : Ridley USA Inc.
111 W. Cherry St., Suite 500, Mankato, MN 56001

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID

If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
If on skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes. Call a poison control center or doctor for treatment advice.

Emergency Assistance: Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), you may call 1-800-222-1222 24 hours per day, 7 days per week.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FEEDING AND MANAGEMENT INSTRUCTIONS:

Provide Prairie Pride Feeds® ClariFly® Fly Control Block as a free choice source of salt and minerals for your cattle and horses. When used as directed, it will provide sufficient ClariFly® to inhibit fly emergence from the manure of treated cattle and horses. Existing adult flies will not be affected. Start feeding this product early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity. Allow free choice to cattle and horses. Cattle should consume an average of 0.8 oz. of Prairie Pride Feeds® ClariFly® Fly Control Block per 100 lbs of animal body weight per day, which is equivalent to 8 oz. per day for a 1,000-lb cow. Horses should consume an average of 1.2 oz. of this product per 100 lbs of animal body weight per day which is equivalent to 12 oz. per day for a 1,000-lb horse.

To Open: Remove top plastic cover sheet.

1. Assure cattle and horses are not starved for nutrients prior to feeding this block.
2. This product contains added copper. Do not feed to sheep and other species that have a low tolerance to supplemental copper. Do not use in association with feeds containing supplemental selenium.
3. Offer cattle and horses one Prairie Pride Feeds® ClariFly® Fly Control Block for every 3 to 5 head.
4. Remove all other self limiting nutritional supplements.
5. Locate Prairie Pride Feeds® ClariFly® Fly Control Block where cattle and horses congregate (watering, loafing, shade areas).
6. Replenish Prairie Pride Feeds® ClariFly® Fly Control Block on a regular basis as needed.
7. If intake of this block is below 0.8 oz. per 100 lbs animal body weight per day for cattle or 1.2 oz. per 100 lbs. animal body weight per day for horses, increase the number of feeding locations and/or change locations to areas more frequented by cattle and horses. If intake of this block is above 0.8 oz. per 100 lbs animal body weight per day for cattle or 1.2 oz. per 100 lbs animal body weight per day for horses, reduce the number of feeding locations and/or change locations to areas less frequented by cattle and horses.
8. Introduction of this product after adult fly infestation is established may require treatment of cattle and horses with supplemental sprays to reduce the adult fly population coming from outside sources.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store in a cool, dry place. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Completely empty wrapper. Dispose of packaging material in a sanitary landfill or by other procedures approved by state and local authorities.

EPA Reg. No. 89459-7-7698
EPA Est. No. 7698-TX-002^F, 7698-MN-003^B
7698-KY-002^H, 7698-KS-001^J, 7698-AL-001^E
Superscript letter is the first letter in batch number.

65447

0219



3 12190 65447 6
NET WEIGHT: 33.3 LB (15.12 KG)