

**PROTECTS  
OVER 125  
LISTED  
FRUITS AND  
VEGETABLES**



**GARDENTECH**

**Sevin®**

**INSECT KILLER DUST**

**READY TO USE**



**KILLS OVER 150 LISTED INSECTS**

**ON VEGETABLES, FRUITS, ORNAMENTALS & LAWNS**

**PROTECTS EDIBLES FROM  
INSECT DAMAGE**

**STARTS KILLING IMMEDIATELY  
UPON CONTACT**

**ACTIVE INGREDIENT:**

Bifenthrin ..... 0.1%

**OTHER INGREDIENTS:** ..... 99.9%

**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

Si usted no entiende la etiqueta,  
busque a alguien para que se la  
explique a usted en detalle.  
(If you do not understand the  
label, find someone to explain it  
to you in detail.)

**Net Weight: 3 LB (1.36 kg)**

Distributed by: TechPac, LLC,  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339

RT032221AAA

Sevin is a registered trademark of Tessenderlo Kerley, Inc.  
Garden Tech is a registered trademark of Gulfstream Home  
and Garden Inc.

EPA Reg. No. 53893-484-7104  
EPA Est. 67572-6A-1, 67572-6A-2, 69703-6A-1, 69703-6A-2,  
Subscription is first letter of main code on container.

For Consumer Return:  
TechPac LLC, PO Box 95437  
Palatine, IL 60095-0437

If you have a medical emergency, need consultation on diagnosis, therapy and analysis, please call  
1-(800)-420-9347. If you have questions relating to product use or application, please call 1-(866)-945-5033.  
If you have a product spill or transportation emergency, please call 1-(800)-424-9300.



Read the label before applying!  
Kills by contact  
Simply shake to control listed insects

People and pets can return to the treated area once the dust has settled

## FIRST AID

|                                |   |
|--------------------------------|---|
| <b>IF SWALLOWED:</b>           | <ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>IF IN EYES:</b>             | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>                                 |
| <b>IF ON SKIN OR CLOTHING:</b> | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |
| <b>IF INHALED:</b>             | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>                             |

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-420-9347 for emergency treatment information.

**NOTE TO PHYSICIAN:** This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and other aquatic organisms. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.

**⚠ NOTICE:** Cancer - [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**IMPORTANT: READ ENTIRE LABEL BEFORE EACH USE.**

### For Outdoor Residential Use Only

This product is a ready-to-use insecticide dust, which exhibits effective control against the pests listed when used according to label directions. This product is non-injurious to most plants; even so, all varieties of plants have not been fully tested. Before treating large numbers of plants of a single variety or first treating material not previously tested, treat a small area and observe for injury prior to full scale application. Wait 5-7 days after treatment to evaluate results. Plants under stress (such as heat, drought, or frost damage) may be sensitive to this product. Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

For most effective results, follow directions for specific use areas.

**WHERE TO APPLY:** Around Home Foundations, Lawns, Shrubs, Ornamental and Flower Gardens, Fruit and Vegetable Gardens.

**WHEN TO APPLY:** Apply early to prevent infestation. Otherwise, apply at the first signs of insect activity or damage.

### RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dust has settled.
- Do not apply to pets or pet bedding/quarters.
- Do not make applications during rain. Avoid making applications when rainfall is expected.
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- Do not apply when the wind speed is greater than 15mph.
- Not for use on plants grown for sale or commercial use, research, or commercial seed production.
- For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Do not allow the product to enter any drain during or after application.
- Do not apply or irrigate to point of runoff.

In New York State, this product may NOT be applied to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

All outdoor applications must be limited to spot or crack-and crevice treatment only, except for the following permitted uses:

1. Treatment to soil or vegetation around structures;
2. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
3. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

## **Around Homes**

**HOW TO APPLY:** To help prevent the invasion of buildings and structures by listed pests, lightly dust a band 3 to 4 inches wide around the outside foundation wall of the home, at a rate of 8 ounces per 1000 ft<sup>2</sup>. Do not apply more than 4.5 pounds per 1000 ft<sup>2</sup> per year.

## **On Lawns**

**HOW TO APPLY:** Apply when insects or their damage first appear and repeat at weekly intervals or as necessary. Apply 8 ounces per 1000 ft<sup>2</sup>. Sprinkle a thin uniform layer of dust on grass. Do not apply more than 4.5 pounds per 1000 ft<sup>2</sup> per year.

## **On Ornamentals & Flowers**

**HOW TO APPLY:** Apply when insects or their damage first appear and repeat at weekly intervals or as necessary. Apply 8 ounces per 1000 ft<sup>2</sup>; or thoroughly apply to place a thin uniform layer of dust on both upper and lower sides of leaves, stems and flowers. Do not apply more than 4.5 pounds per 1000 ft<sup>2</sup> per year.

## **Pest List for Lawns, Around Homes and on Ornamentals & Flowers**

Adelphocoris Plant Bugs, Amorbis spp., Ants (except Fire Ants, Harvester, and Pharaoh), Arizona Cricket, Artichoke Plume Moth, Asian Lady Beetle, Billbugs, Blister Beetles, Boxelder bugs, Boxwood Leafminers, Carpenter Ants (Eastern), Carpenter Ants (Western), Clearwing Moth, Common Short-tailed Cricket, Dark Mealworms, Darkling Beetle, Deer ticks, Earwigs, Eastern Striped Cricket, Elm Bark Beetle, Elm Leaf Beetle, Elm Leaf Rollers, Fall Field Cricket, Firebrat, Flat Grain Beetles, Four-lined Plant Bug, Fuller Rose Beetle, Garden symphylan, German Cockroach, Glassy-winged sharpshooter, Green Fruit Beetle, Holly Leafminers, Hoplia beetle, House Cricket, Indies Short-tailed Cricket, Katydid, Lace Bugs, Leafhopper, Lesser Grain Borers, Lycaenid pod borer, Mealybugs, Mexican Bean Beetle, Millipedes, Mormon Cricket, Northern Wood Cricket, Oak Leafminers, Periodical Cicada, Pillbugs, Potato Beetle, Raspberry Crane Maggot, Raspberry Crown Borer, Raspberry Horntail, Redhumped Caterpillar, Rose Chafer, Rose Midges, Sand Field Cricket, Scarlet Plant Bugs, Scarlet-and-Green Leafhopper, Seedcorn maggot, Sharpshooter, Silverfish, Sod Webworm, Southeastern Field Cricket, Southern Wood Cricket, Sowbugs, Spotted wing drosophila, Spring Field Cricket, Spruce budworm, Spruce Needle Leafminers, Strawberry Crown moth, Tent caterpillars, Texas Field Cricket, Tropical Cricket, Variable Field Cricket, Vocal Field Cricket, Western Striped Cricket, Western Tent Caterpillar Moth

## **Home Garden Vegetables And Small Fruits**

**HOW TO APPLY:** Apply when insects or their damage first appear. Apply 1 lb (16 ounces) per 500 square feet; or thoroughly apply to place a thin, uniform layer of dust on both upper and lower sides of leaves, stems and flowers. Reapply as necessary but do not apply more than 4.5 pounds of product per 1000 ft<sup>2</sup> per year. Observe the following crop-specific use directions, including the preharvest interval (PHI) which is the number of days between last application and harvest.

| <b>GARDEN CROP</b>   | <b>INSECT PESTS</b>   | <b>COMMENTS</b>  | <b>PHI</b> |
|--|---|--|------------|
| <b>Strawberries</b>  | Aphids, Armyworms, Flea Beetles, Corn Earworm (Heliiothispp.), Leafrollers, Lygus Bug (Lygus spp.), Plant Bugs, Spittlebugs, Stink Bugs, Strawberry Clipper Weevil, Strawberry Sap Beetle   | Apply when pest populations reach damaging thresholds and repeat as necessary at 7-14 day intervals. | 0          |
| <b>Caneberries:</b><br>Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, Raspberries   | Black Vine Weevil, Leafrollers, Orange Tortrix, Root Weevils  | One application may be made pre-bloom and a second application may be made post bloom.               | 3          |
| <b>Pears</b>   | Aphids, Codling Moth, Cutworms, Green Fruitworm, Leafhoppers, Leafminers, Leafrollers, Lygus Bug (Lygus spp.), Plant Bugs, Plum Curculio, San Jose Scale (Crawler), Stink Bugs, Tarnished Plant Bugs  | Apply as necessary to maintain control using a minimum of 30-day interval.                           | 14         |
| <b>Succulent Peas and Beans:</b><br>Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea, Pigeon pea, Broadbean (succulent), Lima bean (green), Runner bean, Green bean, Snap bean, Wax bean, Asparagus bean, Blackeyed pea, Chinese longbean, Cowpea, Moth bean, Southern pea, Yardlong bean, Jackbean, Sword bean   | Alfalfa Caterpillar, Aphids, Aster Leafhopper, Bean Leaf Beetle, Beet Armyworm, Cloverworm, Corn Earworm, Corn Rootworm, Cucumber Beetles, Cutworms, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshoppers, Japanese Beetle, Leafhoppers, Loopers, Pea Leaf Weevil, Pea Weevil, Plant Bug, Sap Beetle, Southern Armyworm, Stink Bugs, Tarnished Plant Bug, Thrips, Webworms, Western Bean Cutworm, Yellowstriped Armyworm | Do not make applications less than 7 days apart.   | 3          |
| <b>Head and Stem Brassica Vegetables:</b><br>Broccoli, Chinese Broccoli (Gailon, White Flowering Broccoli), Brussels Sprouts, Cauliflower, Cavalo broccolo, Kohlrabi, Cabbage, Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy)  | Aphids, Armyworms, Corn Earworm, Crickets, Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles, Ground Beetles, Imported Cabbageworm, Saltmarsh Caterpillar, Leafhoppers, Loopers, Stink Bugs, Thrips, Tobacco Budworm, Wireworm   | Do not make applications less than 7 days apart. Do not make more than 5 applications after bloom.   | 7          |
| <b>Cucurbits:</b><br>Chayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Edible Gourd (includes hyotan, cucuzza), Luffa spp. (includes hechima, Chinese okra), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or Cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and Snake melon), Pumpkin, Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter Squash (includes Butternut squash, calabaza, hubbard squash (C. mixta; C. pepo) includes | Aphids, Armyworms, Cabbage Looper, Cutworms, Cucumber Beetles, Corn Earworm, Grasshopper, Leafhoppers, Melonworm, Pickleworm, Plant Bug, Rindworm, Squash Bugs, Squash Vine Borer, Stink Bugs, Tobacco Budworm  | Do not make applications less than 7 days apart. Do not make more than 2 applications after bloom.   | 3          |

| GARDEN CROP   | INSECT PESTS  | COMMENTS  | PHI |
|---|---|---|-----|
| <b>Cucurbits: - continued</b><br>acorn squash, spaghetti squash),<br>Watermelon (includes hybrids<br>and/or varieties of Citrulluspp.).   |   |   |     |
| <b>Lettuce (Head)</b>   | Aphids, Armyworms, Corn<br>Earworm, Cucumber Beetles,<br>Cutworms, Diamondback Moth,<br>Flea Beetles, Imported<br>Cabbageworm, Leafhoppers,<br>Loopers, Salt Marsh Caterpillar,<br>Stink bug, Tobacco Budworm   | Do not make<br>applications less<br>than 7 days apart.  | 7   |
| <b>Soybeans/Edamame</b>   | Soybean Aphid   | Apply foliar<br>treatments at a<br>minimum of 30<br>days intervals.   | 18  |
| <b>Fruiting vegetables (except<br/>Tomato &amp; Tomatillo):</b><br>Eggplant, Pepper (Bell &<br>Non-Bell), Groundcherry, Pepino  | Armyworms, Cabbage Looper,<br>Colorado Potato Beetle, Corn<br>Earworm, Cucumber Beetle,<br>Cutworms, European Corn<br>Borer, Flea Beetle, Leafminers,<br>Loopers, Pepper Weevil, Plant<br>Bug, Stink Bug, Thrips, Tomato<br>Pinworm, Tomato Hornworm  | Do not make<br>applications less<br>than 7 days apart.  | 7   |
| <b>Tomato &amp; Tomatillo</b>   | Aphids, Armyworms, Bean Leaf<br>Beetle, Cabbageworm,<br>Cloverworm, Corn Earworm,<br>Corn Rootworm, Cucumber<br>Beetles, Cutworms,<br>Diamondback Moth, European<br>Corn Borer, Flea Beetles, Flea<br>Hopper, Grasshopper, Japanese<br>Beetle, Leafhoppers, Loopers,<br>Lygus Bug (Lyguspp.),<br>Melonworm, Pea Weevil, Pea<br>Leaf Weevil, Pickleworm, Plant<br>Bug, Rindworm, Salt Marsh<br>Caterpillar, Sap Beetle, Seedpod<br>Weevil, Squash Bugs, Stink bug<br>spp., Tobacco Budworm,<br>Tarnished Plant Bug, Thrips,<br>Wireworms | Do not make<br>applications less<br>than 10 days<br>apart. Do not<br>apply more than<br>4 applications<br>(0.32 lb active<br>ingredient) per<br>season. | 1   |
| <b>Peanut</b>   | Beet Armyworm, Corn Earworm,<br>Cutworm spp., Fall Armyworm,<br>Grasshoppers, Green<br>Cloverworm, Leafhoppers,<br>Lesser Cornstalk Borer,<br>Loopers, Rednecked Peanut<br>Worm, Southern Armyworm,<br>Southern Corn Rootworm, Stink<br>bugs, Three Cornered Alfalfa<br>Hopper, Velvetbean Caterpillar,<br>Yellowstriped Armyworm   | At a minimum of<br>14 days intervals.   | 14  |
| <b>Root Crops (except Sugar<br/>Beets &amp; Garden Beets):</b><br>Burdock (edible), Carrot,<br>Celeriac, Chervil (Turnip Rooted),<br>Chicory, Ginseng, Horseradish,<br>Parsley (Turnip Rooted), Parsnip,<br>Radicchio, Radish (Oriental),<br>Rutabaga, Salsify, Salsify<br>(Black), Salsify (Spanish),<br>Skirret, Turnip | Aphids, Beet Armyworm, Celery<br>Leaf Tier, Corn Earworm,<br>Cross-Striped Cabbageworm,<br>Cutworms, Diamondback Moth,<br>European Corn Borer, Fall<br>Armyworm, Flea Beetles, Green<br>Cloverworm, Hornworms,<br>Imported Cabbageworm,<br>Loopers, Southern Armyworm,<br>Tobacco Budworm, Velvetbean<br>Caterpillar, Yellowstriped<br>Armyworm   | Do not make<br>applications less<br>than 7 days apart.  | 21  |
| <b>Leafy Petiole Vegetables:</b><br>Celery, Cardoon, Chinese Celery,<br>Celtuce, Florence Fennel,<br>Rhubarb, Swiss Chard   | Aphids, Armyworms, Cutworms,<br>Corn Earworm, Crickets,<br>Cucumber Beetles,<br>Diamondback Moth, Flea<br>Beetles, Ground Beetles,<br>Imported Cabbageworm,<br>Leafhoppers, Loopers,<br>Saltmarsh Caterpillar, Stink<br>Bugs, Thrips, Tobacco<br>Budworm, Wireworm  | Do not make<br>applications less<br>than 7 days apart.  | 7   |
| <b>Bushberries:</b><br>Blueberry (Highbush and<br>Lowbush), Currant Elderberry,<br>Gooseberry, Huckleberry  | Aphids, Blueberry Maggot,<br>Fruitworms, Leafhoppers,<br>Lecanium Scale (crawlers),<br>Plum Curculio, Oblique Banded<br>Leaf Roller, Red Banded<br>Leafroller, Spanworm,<br>Variegated Leafroller   | Do not make<br>applications less<br>than 7 days apart.  | 1   |

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in tightly closed container in a dry place.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency or 1-866-945-5033 for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** Read the entire Directions For Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

## WARRANTY STATEMENT

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, User agrees to hold TechPac LLC and Seller harmless for any claims relating to such factors.

