

GOODWINOL PRODUCTS CORP.

P.O. BOX 407, PIERCE, CO, 80650

Telephone: 970-834-1229

Order Desk: 800-554-1080

Fax: 970-834-0180

Website: www.goodwinol.com

Email: goodwinol@what-wire.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

VETR_xTM POULTRY REMEDY

Goodwinol

Active Ingredient(s): Made with 3.3% (v-v) alcohol U.S.P. The mixture contains Canada balsam, camphor, oil origanum, oil rosemary, blended in a corn oil base.

Indications: For use as an aid in the treatment of respiratory problems in all varieties of poultry, including bantams, ducks, turkeys, geese, and game birds.

Effective reliefs for colds, roup, scaly legs, and eye worm and as a conditioner for show preparations. For internal and external application.

Dosage and Administration: Dosages and Preparing Solution: The directions below apply to bantams and mature poultry of similar size, unless otherwise noted. For larger or smaller birds, change the dosage in proportion to the size of the bird.

Use VETR_xTM warm, unless otherwise specified. To warm, place the spout in an open position. Put the bottle in a small pan of water. Heat at medium temperature. Always test the temperature of the product before applying internally or externally.

To prepare the solution, add one (1) teaspoonful of warm VETR_xTM to one (1) cup of very warm water.

VETR_xTM mixes with hot water, but floats on top of cold water.

Colds and Roup: Symptoms of colds are watery eyes, droopy appearance, and an unusual tendency to shake or turn the head.

To treat colds, drip VETR_xTM solution into the bird's nostrils. Place a few drops of warm VETR_xTM, straight from the bottle, down the throat of the infected bird at night. Also, rub some warm VETR_xTM from the bottle over the head of birds and under wings. If a cold develops into more severe roup, change to a stronger solution of one (1) teaspoonful of VETR_xTM in one-half (1/2) cup of hot water. Treat the same as for colds. Also, swab the throat and wash the head and eyes with the stronger solution four (4) or five (5) times a day. Keep the nostrils, eyes, and throat as clean as possible.

Flock Care: If colds infect many in a flock, spray warm VETR_xTM solution over as many birds as possible. Use any small sprayer, and keep the sprayer in a pan of hot water before using. For birds in a flock which look droopy, rub warm VETR_xTM, full strength, over their heads, eyes, and under their wings. Also, provide plenty of clean, cool drinking water for the flock. Add a few drops of VETR_xTM to the drinking water each time it is changed. It will float on the water and get on the beaks and nostrils of the birds each time they drink.

Vaporizer: If deciding to use a vaporizer, make certain that it is a hot water model. Follow the manufacturer's directions. Fill the cup with VETR_xTM full strength and operate the vaporizer for at least one (1) hour while birds are roosting.

Preventing Colds and Roup: To help prevent colds, sprinkle a few drops of VETR_xTM in the litter every four (4) or five (5) days. Add a few drips to the drinking water every time it is changed. Rub one (1) or two (2) drops of warm, full-strength VETR_xTM under the wings.

For Show Birds: To protect birds from colds while traveling to and from shows, place a few drops of warm VETR_xTM under one (1) wing. It does not stain, or harm feathers, wing, or color. Rubbing VETR_xTM on the legs brings out the true color and helps to protect them from mites and lice. When the birds return from

shows, keep them separated from the flock for one (1) week. Wash their feet and heads with a warm solution, and rub warm VETR_x[™] from the bottle on the heads and under wings.

Scaly Legs: When shanks of the bird appear dry, rub them with a small amount of cold VETR_x[™] from the bottle. In cases of heavy scale, apply cold VETR_x[™] and rub in thoroughly all over the shanks and toes. Repeat every three (3) days until the legs are completely free of scale.

“Eye Worm”: The tiny worm works its way under the third eyelid and causes a pus pocket to form. The bird will blink excessively and rub its eye at the base of the wing feather. To treat, hold the bird under one (1) arm so that both hands are free. Open the bird's mouth while turning its head to the side and tilting down to the cleft in the roof of the mouth can be seen. Dip a small cotton swab into warm VETR_x[™]. Press the swab right into the cleft. Do this slowly. Watch for VETR_x[™] to come from each side of the bird's beak and pus coming from the corner of each eye. Remove the pus and rest the bird for awhile. The treatment may have to be repeated two (2) or three (3) times before results are seen.

Tonic Feeder for Breeding: To keep breeding birds healthy for maximum production of breeding eggs. Keep three (3) quarts of laying mash indoors for a few hours until it reaches room temperature. Warm a bottle of VETR_x[™]. Pour the entire bottle onto the mash and mix well by rubbing the mash between the hands. Let it stand in a warm room over night. Next day, put the treated mash in a container with a tight lid and let it stand for about two (2) weeks. Mix one (1) tablespoon of treated mash, once a day, in feed for every 20 birds. If feeding 10 birds or less, one (1) teaspoon is enough. Be sure to stir the treated mash thoroughly into feed.

General Precautions: While treating poultry and other birds, follow the usual precautions. Isolate infected birds as soon as possible. Clean out all coops lofts in which sick birds have been. Keep the birds as clean and dry as possible, and free from drafts. Watch for evidences of disease in birds coming home from shows. Diseases are often contracted in baskets.

If symptoms persist consult a veterinarian.

Caution(s): For animal use only. Not for human consumption. Use only as directed. Keep all medication out of the reach of children.

Presentation: 2 fl. oz. (59 mL) and 1 quart.

CPN: 1438008.0