



GoatVac C.D.-T[®]

Clostridium Perfringens Types C & D Tetanus Toxoid



INDICATIONS:

For use in healthy goats as an aid in the prevention of and enterotoxemia caused by *Clostridium perfringens* types B, C, and D. *Cl. Perfringens* Type B is not a significant problem in the United States. For long-term protection against tetanus.

FOR USE ON:

Goats

PRODUCT FEATURES:

- Safe for all ages
- USDA approved
- High efficacy against disease

CONTAINS:

Thimerosal as a preservative.

DIRECTIONS FOR USE:

Do not vaccinate within 21 days before slaughter. Shake well before use.

Use entire contents when first opened. Store at 2° to 7° C. Do not freeze.

Vaccinate sufficiently in advance of feeding concentrated rations to provide a minimum of 2 weeks after second dose for a adequate immunity to develop.

Tetanus toxoid requires 3 to 4 weeks to establish effective protection that will last several months. Booster injections should be made annually, or, in event of injury, regardless of interval.



PRECAUTIONS:

Anaphylactoid reaction may occur following administration of products of this nature.

If noted, administer adrenalin or equivalent.

DOSAGE and ADMINISTRATION:

Inject 2mL subcutaneously or intramuscularly.

Kids : Initial dose at pre-weaning (30 days of age), Booster at weaning (60 days of age), Annual Booster

Does/Nannies: Initial dose 60 days prior to whelping, Booster 3-4 weeks after initial dose, Booster 6 months after initial dose, Annual Booster

Bucks/Billies: Initial dose, Booster 3-4 weeks after initial dose, Annual Booster

PACKAGING:

20 mL (10 dose) bottle, 10 per case

UPC #: 7 45801 41302 3



DURVET, INC. ANIMAL HEALTH PRODUCTS
P.O. Box 279 · Blue Springs, Missouri 64014 · 800/821-5570 · 816/229-9101
Fax 816/224-3080 · E-mail: info@durvet.com · www.durvet.com