## **VISION® 7 WITH SPUR®\***

## Intervet/Schering-Plough Animal Health

**Cattle Vaccine** 

## CLOSTRIDIUM CHAUVOEI - SEPTICUM - NOVYI - SORDELLII - PERFRINGENS TYPES C & D BACTERIN - TOXOID

For use in healthy cattle and sheep as an aid in preventing disease caused by *Clostridium chauvoei* (Blackleg), *septicum* (Malignant edema), *novyi* (Black disease), *sordellii* and *perfringens* Types C & D (Enterotoxemia).

**DOSE:** Cattle-2mL, Sheep-2mL. Inject subcutaneously. Repeat in 3 to 4 weeks. Annual revaccination is recommended. Shake well before using. Store at 35° to 45°F (2° to 7°C). Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

**ANTIDOTE:** Epinephrine. Contains formaldehyde as preservative.

\*Adjuvant-Intervet's Proprietary Technology

FOR USE IN ANIMALS ONLY

Intervet Inc., Millsboro, DE 19966

1-800-441-8272

www.intervetusa.com

U.S. Vet. Lic. No. 286

			Code	
	10 Doses	20 mL	069130	09581005
	50 Doses	100 mL	069131	09585004
	250 Doses	500 mL	069132	095825004

**CPN:** 1106216.2