INTERVET/SCHERING-PLOUGH ANIMAL HEALTH, Intervet Inc. (See Merck Animal Health)

Distributed by MERCK ANIMAL HEALTH



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Inactive - Content may be out of date

This product has been replaced with 20/20 Vision® 7 with Spur® (Intervet/Merck Animal Health)

20/20 VISION® 7 WITH SPUR®*

Intervet/Schering-Plough Animal Health

Cattle Vaccine

CLOSTRIDIUM CHAUVOEI - SEPTICUM - NOVYI - SORDELLII - PERFRINGENS TYPES C & D - MORAXELLA BOVIS BACTERIN - TOXOID

For use in healthy cattle as an aid in preventing disease caused by *Clostridium* chauvoei (Blackleg), septicum (Malignant edema), novyi (Black disease), sordellii and perfringens Types C & D (Enterotoxemia), and *Moraxella bovis* (Pinkeye, or infectious bovine keratoconjunctivitis).

DOSE: Cattle-2mL. Inject subcutaneously. Repeat in 3 to 4 weeks. Shake well before using. Store at 35° to 45°F (2° to 7°C). Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

ANTIDOTE: Epinephrine. Contains formaldehyde as preservative.

*Adjuvant-Intervet's Proprietary Technology.

FOR USE IN ANIMALS ONLY

Intervet Inc., Millsboro, DE 19966

1-800-441-8272

www.intervetusa.com

U.S. Vet. Lic. No. 286

			Code	
	10 Doses	20 mL	069083	09701003
	50 Doses	100 mL	069084	09705003

CPN: 1106001.2