

(ivermectin) POUR-ON for Cattle
Contains 5 mg ivermectin per mL

IVERMECTIN

durvet

IVERMECTIN



(ivermectin)
POUR-ON
for Cattle

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

INDICATIONS:

For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle.

See package insert for complete indications and use directions.

DOSAGE:

The dose rate is 1 mL for each 22 lb of body weight. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead.

ADMINISTRATION:

Connect the applicator gun to the back pack as follows:

Attach the open end of the draw-off tubing to the dosing equipment. (Because of the solvents used in the formulation, only the Protector Drench Gun from Instruments Supplies Limited, or equivalent, is recommended. Other applicators may exhibit compatibility problems resulting in locking, incorrect dosage or leakage.)

Replace the shipping cap with the draw-off cap and tighten down. Attach draw-off tubing to the draw-off cap.

Gently prime the applicator gun, checking for leaks.

Follow the manufacturer's directions for adjusting the dose.

When the interval between uses of the applicator gun is expected to exceed 12 hours, disconnect the gun and draw-off tubing from the product container and empty the product from the gun and tubing back into the product container. To prevent removal of special lubricants from the Protector Drench Gun, the gun and tubing must not be washed.

IVERMECTIN



(ivermectin)
POUR-ON
for Cattle

ANIMAL SAFETY: Studies conducted in the U.S.A. have demonstrated the safety margin for ivermectin. Based on plasma levels, the topically applied formulation is expected to be at least as well tolerated by breeding animals as is the subcutaneous formulation which had no effect on breeding performance.

RESIDUE WARNING: Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING!
NOT FOR USE IN HUMANS**

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin. To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

Keep this and all drugs out of the reach of children.

TAKE TIME  OBSERVE LABEL DIRECTIONS

**WARNING! FLAMMABLE!
KEEP AWAY FROM HEAT,
SPARKS, OPEN FLAME, AND
OTHER SOURCES OF IGNITION.**

durvet

IVERMECTIN



(ivermectin)
POUR-ON
for Cattle

Contains 5 mg ivermectin per mL

Parasiticide

Kills: Roundworms (including Brown Stomach Worm), Lungworms, Grubs, Sucking Lice, Biting Lice, Mange Mites, Horn Flies

Restricted Drug (California) – Use Only as Directed

Contains 200-550 lb Doses

ANADA 200-318, Approved by FDA

Net Contents: 169 fl oz (5 Liter)

Manufactured for:
Durvet, Inc.
Blue Springs, MO 64014
www.durvet.com

IVERMECTIN



(ivermectin)
POUR-ON
for Cattle

This topically applied formulation of IVERMECTIN (ivermectin) Pour-On delivers internal and external parasite control in one convenient low-volume application.

PRECAUTIONS:

Store at 20°C - 25°C (68°F - 77°F), with the excursions information included on the package insert and protect from light.

Use only in well-ventilated areas or outdoors.

Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced.

Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally.

Cloudiness in the formulation may occur when IVERMECTIN (ivermectin) Pour-On is stored at temperatures below 32°F. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g., caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, IVERMECTIN (ivermectin) Pour-On is not recommended for use in species other than cattle.

Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

Restricted Drug (California) - Use Only as Directed

Product No. 1IVE014
8IVE024 Rev: 01/13



Manufactured for:
Durvet, Inc.
Blue Springs, MO 64014
www.durvet.com