

VISION® 8 WITH SPUR®*

Intervet/Schering-Plough Animal Health

Cattle Vaccine

CLOSTRIDIUM CHAUVOEI - SEPTICUM - HAEMOLYTICUM - NOVYI - SORDELLII - PERFRINGENS TYPES C & D BACTERIN - TOXOID

For use in healthy cattle and sheep as an aid in preventing disease caused by *Clostridium chauvoei* (Blackleg), *septicum* (Malignant edema), *haemolyticum* (Bacillary Hemoglobinuria/Red Water), *novyi* (Black disease), *sordellii* and *perfringens* Types C & D (Enterotoxemia).

DOSE: Cattle - 2mL, Sheep - 2mL. Inject subcutaneously. Repeat in 3 to 4 weeks. In animals subject to reexposure to *Cl. haemolyticum* revaccinate every 5 to 6 months. Annual revaccination is recommended. Shake well before using. Store at 35° to 45°F (2° to 7°C). Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

ANTIDOTE: Epinephrine. Contains formaldehyde as preservative.

*Adjuvant-Intervet's Proprietary Technology.

FOR USE IN ANIMALS ONLY

Intervet Inc., Millsboro, DE 19966

1-800-441-8272

www.intervetusa.com

U.S. Vet. Lic. No. 286

		Code	
10 Doses	20 mL	069136	09591003
50 Doses	100 mL	069137	09595003
250 Doses	500 mL	069138	095925003

CPN: 1106218.2