

SpectoGard® Scour-Check®

(spectinomycin)

ORAL SOLUTION FOR PIG SCOURS
Anti-Infective

ANADA 200-364, Approved by FDA



DESCRIPTION

SpectoGard® Scour-Check® is an oral solution intended as an anti-infective for young pigs. Each mL contains spectinomycin dihydrochloride pentahydrate equivalent to 50 mg of spectinomycin activity.

INDICATIONS

For use in pigs under four (4) weeks of age for treatment and control of porcine enteric colibacillosis (scours) caused by *E. coli* susceptible to spectinomycin.

BENEFITS

- **Effective:** Proven spectinomycin activity
- **Convenient:** No mixing, ready-to-use
- **Palatable:** Readily accepted
- **Fast acting:** Positive results in 3 to 5 days
- **Safe:** Approved by FDA



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1SPE022	240 mL refillable pump bottles	12
1SPE024	1000 mL, resealable plastic bottles	12
1SPE026	1 gallon, resealable	4

WARNINGS

This product is only intended for use in pigs under four (4) weeks of age or weighing less than 15 lbs. Do not administer within 21 days of slaughter.

See reverse side for Administration and Dosage.

Bimeda, Inc.
One Tower Lane, Suite 2250
Oakbrook Terrace, IL 60181 USA
Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035
Email: sales@bimedaus.com



SpectoGard® Scour-Chek® (spectinomycin)

ORAL SOLUTION FOR PIG SCOURS
Anti-Infective

ANADA 200-364, Approved by FDA

**FOR ANIMAL USE ONLY
NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

Restricted Drug (California) – Use only as directed.

INGREDIENTS:

Each mL contains:

Spectinomycin (from spectinomycin dihydrochloride pentahydrate)..... 50 mg

INDICATIONS:

For use in pigs under four (4) weeks of age for treatment and control of porcine enteric colibacillosis (scours) caused by *E. coli* susceptible to spectinomycin.

DOSAGE AND ADMINISTRATION:

See carton containing 240 mL bottle of SpectoGard Scour-Chek for detailed instructions. Each pump of the plunger delivers 1 mL of solution containing 50 mg of spectinomycin.

Pigs under 10 lbs: 1 pump (1 mL) twice daily.

Pigs over 10 lbs: 2 pumps (2 mL) twice daily.

Treatment may be continued twice daily for 3 to 5 days. If pigs do not improve within 48 hours, rediagnosis is suggested.

DIRECTIONS FOR USE:

The plastic doser supplied in each package with SpectoGard® Scour-Chek® makes treatment easy. After inserting the doser in the bottle, screw the top on tightly to avoid spilling. Push the piece of clear plastic tubing over the end of the pump. Then press the plunger a few times to fill the pump and the clear plastic tube with medication. To administer SpectoGard® Scour-Chek®, insert the plastic tube in the pig’s mouth and press the plunger to obtain the desired dose.

When not in use remove the plunger from the bottle. Put the clear plastic tube in the neck of the bottle and press the plunger a few times to remove the medication from the pump. Replace the original cap on the bottle and always rinse out the doser with water to prevent sticking.

The 1000 mL bottle and gallon jug of SpectoGard® Scour-Chek® should be used only to refill the 240 mL container in the following manner.

1. Remove the cap and doser from the 240 mL bottle or gallon jug.
2. Unscrew the cap from the 1000 mL bottle or gallon jug.
3. Pour, from the 1000 mL bottle or gallon jug, the volume desired into the 240 mL refillable bottle.
4. Reinsert the doser in the 240 mL bottle and screw the top on tightly.
5. Screw the top tightly on the 1000 mL bottle or the gallon jug. Follow the detailed instructions for use shown on the carton containing the 240 mL bottle of SpectoGard® Scour-Chek®.

STORAGE:

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

WARNING:

This product is only intended for use in pigs under four (4) weeks of age or weighing less than 15 lbs. Do not administer within 21 days of slaughter.

HOW SUPPLIED:

SpectoGard® Scour-Chek® is supplied in 240 mL refillable plastic pump bottles, 1000 mL resealable refills and one gallon plastic jugs.

To obtain an MSDS or for assistance, contact Bimeda, Inc. at 1-888-524-6332.

