

FOR USE IN ANIMALS ONLY

INDICATIONS: To be used as an aid in the treatment of parturient paresis (milk fever) in cattle.

DOSAGE AND ADMINISTRATION:

Cattle: 250 to 500 mL

Administer intravenously or intraperitoneally.

Solution should be warmed to body temperature and administered slowly. This product is sterile in unopened container. It contains no preservative. Use entire contents when first opened. It should be handled under aseptic conditions. If there is no apparent improvement in the animal's condition within 24 hours, consult a veterinarian.

TAKE TIME  OBSERVE LABEL DIRECTIONS



ISS19XB04

Manufactured for:
DURVET, INC.

Blue Springs, Missouri 64014
www.durvet.com



NDC 30798-126-17



Calcium Gluconate

23% SOLUTION

Dura-Ster TS

TERMINALLY STERILIZED



**NET
CONTENTS:
500 mL
(16.9 fl. oz.)**

KEEP OUT OF REACH OF CHILDREN

EACH 100 mL CONTAINS:

ACTIVE INGREDIENTS

Calcium Gluconate
(provides 2.14 grams Calcium) 23 grams

INACTIVE INGREDIENTS

Water for Injection q.s.
Boric Acid
(as a stabilizing agent) 3.5 grams
Electrolytes per 1000 mL:
Calcium 1069 mEq
Hypertonic pH 2.5-4.5

Store between 15°C and 30°C (59°F-86°F).