

TRIANGLE® 4 + TYPE II BVD

Boehringer Ingelheim

Bovine Rhinotracheitis-Virus Diarrhea-Parainfluenza-3-Respiratory Syncytial Virus Vaccine

Killed Virus

For Veterinary Use Only

For vaccination of healthy cattle as an aid in the prevention of disease caused by infectious bovine rhinotracheitis (IBR), bovine viral diarrhea (BVD types I & II), parainfluenza-3 (PI-3) and bovine respiratory syncytial (BRSV) viruses. This product contains BVD Types I and II.

DOSE: Cattle, inject one 2 mL dose intramuscularly or subcutaneously using aseptic technique. Repeat in 14 to 28 days. A 2 mL booster dose is recommended annually or prior to time of stress or exposure. Calves vaccinated under six months of age should be revaccinated at six months of age. May be administered to pregnant animals at any stage of gestation. Protect animals from exposure for at least 14 days after the last dose of vaccine.

CAUTION: Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Transient swelling may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine.

Thimerosal, neomycin, and polymyxin B added as preservatives.

© 2007 Fort Dodge Animal Health. All Rights Reserved.

Fort Dodge Animal Health, Fort Dodge, Iowa 50501 USA

U.S. Vet. License No. 112

| | | |
|--------|----------|-------|
| 20 mL | 10 Doses | 1785D |
| 100 mL | 50 Doses | 1787D |

CPN: 1028225.0