This Product Information Sheet is not intended to replace the Product Label. Always read the Product Label for the most current nutrient content and feeding directions.

# 30-15

**Product number:** 8646




## Product Information

- Unique combination of essential nutrients help promote better forage utilization
- Ideal for cattle on low quality forages or pastures
- Reliable, controlled intake averaging 0.75 lb. per head daily
- Weather-resistant formula eliminates waste, saving you money
- One labor-saving, 200-lb tub will supplement 20 head for approximately 2 weeks

## Feeding Instructions

**CAUTION:** Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

**WARNING:** This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

**Introductory Period:** Following a 7 to 10 day acclimation period, average consumption should be 0.5 to 1.1 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.

**Feeding Recommendations:**

1. Feed at a rate of one block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume 0.5 to 1.1 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate SWEETLIX ® self-fed mineral supplement in addition to this supplement.

## Nutrition Information

| | |
|---|---|
| Crude Protein, Min | 30.0 |
| Crude Fat, Min | 3.0 |
| Crude Fiber, Max | 3.0 |
| Calcium, Min | 1.0 |
| Calcium, Max | 1.5 |
| Phosphorus, Min | 1.0 |
| Salt, Min | 2.0 |
| Salt, Max | 2.5 |
| Magnesium, Min | 1.0 |
| Cobalt, Min | 10 |

| | |
|---|---|
| Copper, Min | 300 |
| Iodine, Min | 15 |
| Manganese, Min | 1200 |
| Selenium, Max | 6 |
| Zinc, Min | 900 |
| Vitamin A, Min | 70,000 |
| Vitamin D-3, Min | 7,000 |
| Vitamin E, Min | 70 |

## Product Ingredients

Molasses Products, Processed Grain By-Products, Hydrolyzed Feather Meal, Urea, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Vegetable Oil, Magnesium Oxide, Calcium Carbonate, Plant Protein Products, Roughage Products, Manganese Sulfate, Selenium Yeast, Zinc Sulfate, Zinc Proteinate, Copper Sulfate, Manganese Proteinate, Manganous Oxide, Copper Proteinate, Zinc Oxide, Cobalt Proteinate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil, Ethylenediamine Dihydroiodide, Cobalt Carbonate and Calcium Iodate.