

BIOADVANCED
SCIENCE-BASED SOLUTIONS


3 IN 1

WEED & FEED

for SOUTHERN LAWN

- 1 KILLS**
WEEDS* INCLUDING DOLLARWEED, DANDELION & CLOVER
- 2 FEEDS**
GREENS & STRENGTHENS YOUR LAWN
- 3 PREVENTS**
BROADLEAF & GRASSY WEEDS* PLUS CRABGRASS UP TO **6 MONTHS**

TREATS UP TO
8,000 Sq Ft
NET WT 20 Lbs (9 kg)



BEFORE

AFTER

Active Ingredients:
Dicamba acid 0.088%
Penoxsulam 0.041%
Indaziflam 0.027%
Other Ingredients 99.844%

Total 100.000% *LISTED

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional
precautionary statements

35-0-3

FOR USE ON
SOUTHERN TURFS
BERMUDAGRASS, ST AUGUSTINEGRASS,
CENTIPEDEGRASS & ZOYSIA

QUICK FACTS:

Kills broadleaf weeds* including Dollarweed, Dandelion, Clover, Chickweed & Henbit.

Feeds & strengthens the lawn. Contains slow release fertilizer for long lasting feeding.

Prevents grassy and broadleaf weeds* such as Crabgrass, Annual Bluegrass, Common Dandelion, Goosegrass & Spurge up to 6 months.

WHEN TO APPLY:

Apply anytime in early spring or fall.

WHERE TO USE:

Use only on established lawns of the turf types listed below:
Bermudagrass, Centipedegrass, all varieties of St. Augustinegrass (including Floratam) and Zoysia.

Do not use on any other turf types including: Bahiagrass, Bentgrass, Ormond variety of Bermudagrass, Buffalograss, Carpetgrass, Dichondra, Fescue, Kikuyugrass, Ryegrass, or Seashore Paspalum lawns.

COVERAGE:

This bag treats 8,000 sq. ft.

READ THE LABEL FIRST! DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

- Use this product only on established Southern lawns listed on this label – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded.
- Do not apply this product to water-saturated soils.
- Apply only to actively growing turf.
- Apply in spring after the lawn is actively growing and has been mowed 2 times.
- Do not apply while turfgrass is just turning green in early spring.
- Apply in fall no closer than 6 weeks before the first historical frost date for your area.
- Do not apply to Centipedegrass after October 1.
- Do not apply this product if the lawn appears unhealthy e.g., due to lack of water (drought, watering restrictions), turf disease (dollarspot, large patch, grey leaf spot, take-all root rot, etc.), parasitic microscopic worms (nematodes), turf insects (chinch bugs, etc.), heavy shade, recent aeration or verticutting, or if the lawn is prone to winter injury.
- Do not allow granules to come into contact with flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE

- Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules.
- Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn.
- Wait 24 hours after applying then activate residual barrier by lightly watering in.
- Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn.
- Do not make more than two applications per site in any 12 month period.
- Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

WHEN TO APPLY:

- Apply when - weeds are small and actively growing - temperatures are between 50-90° F.
- May be applied to dry or (for best results) to dew-wetted lawn.
- To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80° F range.
- To prevent Annual Bluegrass (Poa annua) apply August - December.
- Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

MOWING:

- If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product.
- Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING:

- If lawn is dry, water 1 to 2 days before using this product.
- Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS

This product contains a proven blend of weed killers to kill existing weeds* plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds* are dead within weeks. Plus, it feeds your lawn for quick greening and long lasting feeding.


Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.




*listed

SPREADER SETTINGS

BROADCAST SPREADER



DROP SPREADER



Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.

- Tip: 1,000 sq. ft. = 20 ft. x 50 ft.
- Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

BROADCAST SPREADER	SETTING
Earthway Ev-N-Spread	15
Sta-Green	5
Scotts 2000	4 1/2
Scotts Mini	4 3/4
Scotts Deluxe Edge Guard	4 3/4
Scotts Edge Guard Pro	4 1/4

SETTING TO DELIVER 2.5lb/1000 sq. ft.

DROP SPREADER	SETTING
Scotts Accugreen 1000	5 3/4
Scotts Accugreen 2000	6 1/2
Scotts Accugreen 3000	4 1/4

If your spreader is not listed, please call our Consumer Service Hotline toll-free 1-(877) 229-3724 for help.

KILLS

Beggarweed, Creeping
Beggarweed, Dwarf
Betony, Florida
Burdock
Buttercup, Hairy
Buttonweed, Virginia¹
Clover, Hop
Clover, Red
Cocklebur, Common
Coffeeweed,
(Common Chickory)
Daisy, English (Lawn Daisy)
Dandelion, False

Dollarweed (Pennywort)
Goldenrod
Hawkweed
Heallal
Ivy, Ground (Creeping Charlie)
Knotweed
Kyllinga, Green[†]
Lespedeza, Common
Mallow, Roundleaf
Matchweed
Nutsedge, Purple[†]
Nutsedge, Yellow[†]
Parsley-piert

Peppergrass
Pepperweed, Virginia
Plantain, Broadleaf
Smartweed
Sorrel, Sheep
Speedwell, Common
Spotted Catsear
Thistle
Vetch, Common
Wild Violet
Woodsorrel, Creeping
[†]Suppression

PREVENTS

Ageratum, Tropic
Barley, Mouse
Barnyardgrass
Bittercress
Black Nightshade, Hairy
Bluegrass, Annual (Poa annua)
Brome, Foxtail
Buckwheat, Wild
Burweed, Lawn
California Burclover
Chamomile, False
Cheatgrass

Crabgrass, Henry
Crabgrass, Large (Hairy Crabgrass)
Crabgrass, Smooth
Cudweed, Linear Leaf (Purple Cudweed)
Dandelion, Cat's Ear
Dowweed
Evening Primrose, Common
Evening Primrose, Cutleaf
Flatsedge, Rice

Fleabane, Hairy
Foxtail, Giant
Foxtail, Green
Foxtail, Yellow
Filaree, Redstem
Goosegrass
Groundsel, Common
Guineagrass
Kochia
Kyllinga, Fragrant
Lovegrass, Tufted
Mallow, Little
Marestail

Mulberry Weed
Mustard, Black
Mustard, Wild
Nightshade, American
Black
Nightshade, Hairy
Parthenium
Phyllanthus, Longstalked
Pigweed, Prostrate
Pigweed, Redroot
Plantain, Paleseed
Puncturevine
Redmaids

Rocket, London
Ryegrass, Italian
Ryegrass, Perennial
Sandbur
Sedge, Annual
Sesbania, Hemp
Sida, Prickly (Teaweed)
Southern Brassbuttons
Southistle, Annual
Speedwell, Corn
Spurge, Prostrate
Sunflower, Common
Swinecress

Tassel Flower
Velvetleaf
Vetch, Purple
Wild Carrot
Willowherb
Woodsorrel
(Oxalis)

KILLS AND PREVENTS

Burnweed, American
Chickweed, Common
Chickweed, Mouseear
Clover, White
Dandelion, Common
Dock, Curly
Eclipta
Henbit
Knotweed, Prostrate
Kyllinga, Annual[†]
Lambquarters

Medic, Black
Plantain, Buckhorn
Purslane, Common
Pusley, Florida
Ragweed, Common
Shepardspurge
Spurge, Spotted
Thistle, Canada
Woodsorrel, Yellow
[†]Suppression

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call toll-free 1-877-229-3763 for medical emergency information.** Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% Indaziflam (CAS # 950782-86-2).

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS

CAUTION: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

- Harmful if swallowed or absorbed through skin.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- Remove and wash contaminated clothing before reuse.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing and wash hands.
- If product gets on skin, wash thoroughly with soap & water.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Ground Water Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

BioAdvanced® Science-Based Solutions
3-in-1 Weed & Feed for Southern Lawns
35-0-3

GUARANTEED ANALYSIS

Total Nitrogen (N) 35.0%
35% Urea Nitrogen[†]

Soluble Potash (K₂O) 3.0%
Sulfur (S) Total 4.6%

3.5% Free Sulfur (S)
1.1% Combined Sulfur (S) **F2414**

Derived From: Urea, Polymer Coated Sulfur
Coated Urea, & Potassium Sulfate

[†]Contains 17.5% slowly available Nitrogen
from Polymer Coated Sulfur Coated Urea

Information regarding the contents and levels of metals in this product are available on the Internet at <http://www.aapfc.org/metals.html>

Check with your county or city government to determine if there are local regulations for fertilizer use.


NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

****Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

Questions or comments, call toll-free 1-(877) 229-3724 or visit us online at www.bioadvanced.com

EPA Reg. No. 92564-61 EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.
(26) = 82757-FL-1 (57) = 071106-GA-003

BioAdvanced® is a registered trademark of SBM Life Science Corp.
Distributed by: SBM Life Science Corp. 1001 Winstead Drive, Suite 500, Cary, NC 27513



400235661 180502V1