



Malathion

50% SPRAY

BROAD-SPECTRUM CONTROL OF LISTED GARDEN INSECTS
PROTECTS VEGETABLES, FLOWERS, AND SHRUBS

ACTIVE INGREDIENT:
 Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate 50%
INERT INGREDIENTS: 50%
 TOTAL 100%

Contains xylene range aromatic petroleum solvent.
 This product contains 4.35 pounds of Malathion per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear (goggles, face shield, safety glasses). Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Immediately call a poison control center or doctor. Do not induce vomiting unless told to by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate. May cause aspiration pneumonia hazard. This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine. Malathion is an organophosphate insecticide/miticide.

Environmental Hazards

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Physical Or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- **DO NOT APPLY THIS PRODUCT IN A WAY THAT WILL CONTACT ANY PERSON OR PET, EITHER DIRECTLY OR THROUGH DRIFT.**
- **KEEP PEOPLE AND PETS OUT OF THE AREA DURING APPLICATION.**
- For outdoor use only.
- Do not allow people or pets to enter the treated area until sprays have dried.
- When applied as a fogger, do not enter treated area until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- Do not reuse measuring utensils for food use.
- **In New York State: Not for use in residential automatic misting systems.**

TABLE 1. Uses, dosage rates, harvest intervals, and comments for the listed foliar application of Malathion 50% Spray.

Site	Pests	Amount In 1 Gallon of Water	Days to Harvest †	Comments
Tomatoes	Spider Mites Aphids	3 tsp	1	The maximum number of applications per year is 4, and the minimum retreatment interval is 5 days.
Beans (Dry & Succulent)	Mexican Bean Beetles, Spider Mites, Leafhoppers	3 tsp	1	The maximum number of applications per year is 2, and the minimum retreatment interval is 7 days.
Peas	Pea Aphids	3 tsp	3	Make one or more foliar applications. Do not use pea vines for forage. The maximum number of applications per year is 2, and the minimum retreatment interval is 7 days.
White Potatoes	Aphids	3 tsp	0	The maximum number of applications per year is 2, and the minimum retreatment interval is 7 days.
Cabbage	Aphids, Imported Cabbageworm	3 tsp	7	Make full cover applications. The maximum number of applications per year is 6, and the minimum retreatment interval is 7 days.
Kale and Mustard	Aphids, Imported Cabbageworm	3 tsp	7	Make full cover applications. The maximum number of applications per year is 3, and the minimum retreatment interval is 5 days.
Broccoli	Cabbageworm	3 tsp	2	Make full cover applications. The maximum number of applications per year is 2, and the minimum retreatment interval is 7 days.

(cont. on next page)

TABLE 1. (cont.) Uses, dosage rates, harvest intervals, and comments for the listed foliar application of Malathion 50% Spray.

Site	Pests	Amount In 1 Gallon of Water	Days to Harvest ¹	Comments
Turnip	Cabbageworm	3 tsp	1	Make full cover applications. For turnip greens, the maximum number of applications per year is 5, and the minimum retreatment interval is 7 days.
			1	For turnip root, the maximum number of applications per year is 7, and the minimum retreatment interval is 7 days.
Ornamental and/or shade trees, ornamental herbaceous plants, ornamental non-flowering plants, ornamental woody shrubs and vines	Oyster Shell Scale, Euonymus Scale	1 tsp	—	Apply in summer when infestation occurs. Repeat in 10 to 12 days.
	Aphids, White Fly, Thrips, Mealy Bugs, Japanese Beetle (Adult), Four Lined Plant Bug, Rose Leafhopper, Potato Leafhopper	1.5 tsp	—	For scales, apply when crawlers have settled on foliage. Injury may occur on Ferns, Hickory, Viburnum, Lantana, Crassula and Canareti Juniper. Slight injury has also been reported on Boston fern, Pteris fern, and Maidenhair fern, Petunia, Small leaf spirea, Spirea, White pine and Maples. The maximum number of applications per year is 2, and the minimum retreatment interval is 10 days.
	Bagworms, Juniper Scale	2 tsp	—	
	Soft Brown Scale	2.5 tsp	—	
	Pine Needle Scale	4 tsp	—	

¹ Days to harvest refers to the interval or the number of days before harvest for the last application of malathion.

FOR LISTED VEGETABLES USING HOSE-END SPRAYERS

Three Steps To Mix And Apply:

1. FILL

- Determine (Measure) the total area of the garden to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- The application rate for this product is 1.0 fl.oz. of product per 1,000 sq.ft. Add an adequate amount of insecticide to the spray bottle to treat the total area, or add 1 fl.oz. of product for every 1,000 sq.ft. of area.
- Do not add water to the sprayer jar.

2. SET THE DIAL

Set the dial to the rate per gallon and keep the dial set to this setting while spraying. **The dial setting or rate per gallon for this product is 1 Tbs or 3 Tsp oz. of product per gallon of water.**

Types of hose-end sprayers	Dial setting
Ortho® Dial'N Spray® Hose-End Sprayer	3 teaspoons or 1 tablespoons
Hudson® Hose End Liquid Concentrate Sprayer	3 teaspoons or 1 tablespoons
Gilmour® Dial-A-Mix® Lawn and Garden Sprayer	3 teaspoons or 1 tablespoons

Equal measures: 1 tablespoon = 3 teaspoons and 1 fluid ounce = 6 teaspoons or 2 tablespoons

3. SPRAY

Apply evenly over the area. Thorough coverage is important. Rinse sprayer with clean water after each use. Dispose the rinsate over the treated area.

FOR LISTED VEGETABLES WITH BACKPACK, HAND-OPERATED, OR HAND-HELD SPRAYERS

Mix 1 fl.oz. of this product per 2.5 gallons of water for treatment of approximately 1,000 sq.ft.

Gallons of spray	Amount of product, fl.oz.	Amount of product, Teaspoons	Treated area, sq.ft.
1 gallon	0.4 fl.oz.	2.5 teaspoons	400 sq.ft.
2 gallons	0.8 fl.oz.	5 teaspoons	800 sq.ft.
2.5 gallons	1.0 fl.oz.	6 teaspoons	1,000 sq.ft.

Cleaning backpack, hand-operated, or hand-held sprayers: Rinse sprayer with water and or with a cleansing solution of water plus a liquid dishwater detergent. Add 1 to 2 tablespoons of dishwater detergent with one gallon of water to the sprayer tank. Pump the cleaning solution through the spray nozzle onto the treated area. Dispose the remaining cleansing solution over the treated area.

FOR ORNAMENTAL AND/OR SHADE TREES, ORNAMENTAL HERBACEOUS PLANTS, ORNAMENTAL NON-FLOWERING PLANTS, ORNAMENTAL WOODY SHRUBS AND VINES USING HOSE-END SPRAYERS

Three Steps To Mix And Apply:

1. FILL

- Pour product into sprayer jar or add 8 to 12 fl.oz. of product to the sprayer jar.
- Do not add water to the sprayer jar.

2. SET THE DIAL

Set the dial to the rate per gallon and keep the dial set to this setting while spraying. **The dial settings or rate per gallon for this product is for the listed pests below:**

Listed Pests	Rate Per Gallon	Dial Settings
Oyster Shell Scale, Euonymus Scale	1 tsp	1 teaspoon
Aphids, White Fly, Thrips, Mealy Bugs, Japanese Beetle (Adult), Four Lined Plant Bug, Rose Leafhopper, Potato Leafhopper	1.5 tsp	1.5 teaspoons
Bagworms, Juniper Scale	2 tsp	2 teaspoons or 1 tablespoons
Soft Brown Scale	2.5 tsp	3 teaspoons or 1 tablespoon
Pine Needle Scale	4 tsp	4 teaspoons

3. SPRAY

Connect to garden hose to sprayer and begin spraying. Thorough coverage is important. After spraying, unused product in the sprayer jar can be returned (poured) to the original container. Rinse sprayer jar with clean water after each use. Dispose the rinsate over the treated area.

MOSQUITOES AND OTHER SMALL FLYING LISTED INSECTS OUTDOORS:

Amount to Use: Mix 11 tablespoons of product per gallon of water, fuel oil, or diesel oil. Treat lower outside foundation of house, fences, and/or spot treat shrubbery and vegetation where mosquitos rest. Repeat as needed. Do not apply oil solutions to ornamentals, injury may occur.

TABLE 2. Outdoor Uses - Mosquitoes and Other Flying Listed Insects Outdoors for Yards, (Spot treatment only).

Site	Pests	Amount Per Gallon ¹	Comments
Yards – (Spot treatment only), Lower foundations of homes	Mosquitoes, Flying Insects (as previously listed):	11 Tablespoons	Apply as a spray. Repeat as needed. Do not apply oil solutions to ornamental species as injury may occur. Apply as spot treatment for yards and along lower outside foundation of home.

¹ Water, fuel oil, or diesel oil may be used as the carrier or diluent.

Gallons of spray	Amount of product, fl.oz.	Amount of product, Tbs
1 gallon	5.5 fl.oz.	11 Tbs
2 gallons	11.0 fl.oz.	22 Tbs
3 gallons	16.0 fl.oz.	32 Tbs

